


Your Neighbor— The California Ground Squirrel

Otospermophilus beecheyi



**Bill Tietje
Extension Specialist**




Brief History



Santa Barbara Mission--1808

First Vertebrate Pest Conference
(Sacramento, 1962)
\$1.00



The Bounty!
(repealed 1877)



Annual Cycle

Hibernate (Sep-Jan)




Active (Feb-Jun)

-----Competition/Control-----



Aestivate (Jun-Aug)



Next slides:

(Reproduction)

(Predators,
Cattle/Squirrel)

(Oaks) END

Reproduction/Sex Ratio




- 1 litter/year
- 7 young nursed underground 6 wks.
- Clusters of females; males disperse
- Jvs 50%; Yrls 30% (reproductively mature)

Predators



- 70% diet rattlesnake
 - Blood proteins



Predict #s: More when dry.



Cattle & Squirrel

A. Forage Competition

- Reduce gain by 1 lb./day
- 200 squirrels = one 1,000 lb. steer
- 3% of what available (range 2-6%)

Most critical time: Late winter (period of inadequate forage growth for cattle.



B. Study: Fehmi, Russo, Bartolome. 2005. *Effects of Livestock on CA Gd Sq.* J. Range Ecology & Mgmt.

Conclusion: At moderate grazing intensity and during years of adequate rainfall, can have cattle, squirrels, & predators the squirrels support.





Oak Tree & Squirrel

- Provides ideal habitat
 - shelter
 - food (acorns)
- Burrowing dries roots
- Soil to surface may be contaminated w/oak root fungus



END