UNDERSTANDING WORKING RANGELANDSGrazing Systems Management

ACHIEVING MANAGEMENT GOALS BY BALANCING LIVESTOCK GRAZING WITH TIME AND SPACE

SHEILA BARRY, University of California Cooperative Extension Livestock and Natural Resources Advisor, San Francisco Bay Area, and County Director, Santa Clara County; STEPHANIE LARSON, University of California **Cooperative Extension Livestock** and Range Management Advisor, Sonoma and Marin Counties, and County Director, Sonoma County; LAWRENCE D. FORD, Rangeland Conservation Science Consultant, Felton, California; and LISA **BUSH**, Rangeland Management Consultant, Sebastopol, California. The earliest grazing systems date back to the domestication of livestock. Nomadic herdsmen moved livestock from one range site to another, probably following the patterns of forage quality and quantity and the availability of water. Grazing systems became more structured, employing fencing and developed water, over 500 years ago in Europe, when human population pressures demanded greater productivity from agricultural land.

Early ranchers in North America herded or turned livestock loose across open range. Without fences and with natural sources of water found only in limited locations, livestock moved from one site to the next based on the availability of forage and water. The invention of barbed wire in the late 1800s led to better control of livestock and the development of grazing systems in North America. Specialized grazing systems, such as rotation of grazing between pastures, were first conceptualized before the turn of the twentieth century (Smith 1895) and became a focus of range researchers and managers



Photo: Lawrence Ford