



UNIVERSITY OF CALIFORNIA

Division of Agriculture and Natural Resources

<http://anrcatalog.ucdavis.edu>



California Rangelands Research and Information Center

<http://agronomy.ucdavis.edu/calrng/range1.htm>



Guidelines for Monitoring the Establishment of Riparian Grazing Systems

T. A. WARD is University of California Cooperative Extension Livestock and Natural Resource Advisor, San Joaquin and Stanislaus Counties; **K. W. TATE** is UCCE Extension Rangeland Watershed Specialist, Department of Agronomy and Range Science, UC Davis; and **E. R. ATWILL** is Associate Veterinarian and Associate Specialist, Department of Population Health and Reproduction, School of Veterinary Medicine, UC Davis.

This monitoring document outlines methods that will assess current riparian conditions and quantify changes in a riparian area under new management. The monitoring plan outlined here is fairly involved and requires some technical expertise, and for that reason this publication is intended for those with technical experience in rangeland management, specifically UC Cooperative Extension (UCCE) advisors, Natural Resources Conservation Service (NRCS), U.S. Forest Service, and Bureau of Land Management staff, and professional rangeland managers. A secondary audience of land owners and managers can benefit from this information if they are willing to invest time and effort into learning the necessary tools.

Appendixes C1 and C3 at the end of the publication are blank forms that you can copy and use for your own data collection. We have also provided filled-out samples of these and other useful forms to give you a better idea of how to use them.

WHY MONITOR?

When establishing a new riparian grazing system, one would like to be able to compare the success of the new system with the old. Such a comparison can provide validation that the “new and improved” management system is positively affecting riparian health and is a successful project, or that more management changes need to be implemented in order to obtain desired goals, it is through the systematic monitoring of specific conditions that a land manager can assemble this kind of information. The decision left to the manager is, “What tools should I use to assess and monitor my riparian area?” There are a number of ways for managers to conduct their own monitoring, but up until now little guidance has been available as to which tools will best show the results of changes in riparian management. The paragraphs that follow provide an outline for monitoring that is based upon published methods that will provide feedback to changes in riparian grazing management.

SHOULD I MONITOR THE SAME THINGS AS MY NEIGHBOR?

There are distinct benefits to using the same monitoring methods on your property as are used on neighboring properties. If a number of land managers were to implement changes to their riparian areas and each were to select a different set of monitoring tools, they would not be able to compare the relative changes in riparian health between their areas. If, on the other hand, they were to use a common set of monitor tools to observe and record changes in the same items at all sites, they would be able to share information and learn from one another's efforts.

HOW SOON WILL I SEE RESULTS?

Some changes in riparian health can be documented in the short term (a few months to a year), depending on the status of the area at the time you implement management changes. Some changes in riparian health will be observed over the long term (2 years or more). For example, within a year there could be an increase in willow growth (short term), but a change in the tree canopy will not occur for many years (long term). Please see [Appendix A](#) for more information.





Figure 1. Overview of design layout. Transects are established on one side of the creek with their starting points 72 feet apart, providing a total of 360 linear feet (a). Transects should not cross each other, but depending on the site they need not be parallel (b). Be sure to make accurate note of the location of both ends of each transect so you will be able to find them again at a later date.

GETTING STARTED

Before starting, please review all of the published protocols and make sure that you receive any necessary training. [Appendix B](#) of this publication is a sequential outline of the steps necessary to complete the monitoring described here. If you do require training, please contact a UCCE, NRCS, or Resource Conservation District (RCD) office for assistance, possibly including assistance in getting the necessary equipment. The methods do require time and effort, especially during the first year when you first establish the transects. The time required can range from a half day to a full day for two people. Two two-person teams can divide the work and complete the monitoring in less time. It is important that you allow adequate time to collect the necessary data.


ESTABLISHING PERMANENT MONITORING TRANSECTS

To successfully document riparian health changes, you need to be able to examine the same geographical points repeatedly over time. This will ensure that apparent changes in riparian health are the actual results of management and not simply the unique conditions peculiar to different sites. You will need to select a representative section of the riparian area for monitoring; a total of 360 linear feet is required. Six transect lines going across the riparian area and spaced 72 feet apart are established perpendicular to the creek and can be marked using a variety of items such as existing fence posts, lengths of rebar, or wooden stakes painted a unique color (see [Figures 1 and 2](#)). When selecting the marker, give particular thought to the way the pasture is used, the marker's visibility, and its likely permanence over time. If you relate the markers to a benchmark (a permanent fixture such as a tree or large rock), it will be easier for you to find the location of missing markers later on. Record the bearing and distance from the benchmark to each marker.


Transects should encompass upland vegetation on both sides of the creek in order to document whether the width of the riparian vegetation area is increasing or decreasing over time. Because of this, transect lengths will vary from site to site. For example, transects for a mountain meadow system may be 200 feet long, whereas for an intermittent creek in the San Joaquin Valley they may be only 50 feet long. Once you have established the transects, you are ready to begin gathering data.

VEGETATION

To characterize the vegetation, use the USDA Forest Service's Greenline protocol. It is a standard system for classifying and characterizing vegetation and is well suited to



a



b

Figure 2. Examples of transects in the field at different sites (not drawn to scale): *a* has a limited riparian area and so requires shorter transects, while the transects in *b* cross the entire meadow.

the kind of work we are discussing. This protocol has been published, and you should make yourself familiar with its methods and seek out assistance and training if necessary. There is one change that you can make to the Greenline protocol to make it easier to use: key the vegetation according to functional groups instead of species to allow for ease of use while still providing documentation of trends in vegetation succession.

Greenline consists of three components. The first component, *vegetation cross-section composition*, provides information on the width of the riparian area. All six cross-sections are considered for this component. The second, *greenline composition*, was developed for perennial mountain meadows, but is useful for other systems. It documents changes in the permanent greenline along the stream. For example, annual systems may consist of oak trees as the permanent green vegetation. One would expect perennial grasses and other woody species to increase along the stream as management changes were implemented, thus providing a new greenline. The third component, *woody species regeneration*, accounts for any increases in willows, aspen, alders, or other woody plants that tend to provide more stability and canopy cover for the stream. The latter two components are conducted along the permanent vegetation areas of Transect 1 to 6 on both sides of the stream.

VISUAL ASSESSMENTS

Visual assessments are valuable for providing a quick examination of the habitat and hydrologic condition of a system. We recommend that you use two assessments: the U.S. Department of the Interior Bureau of Land Management's *Proper Functioning Condition (PFC)* and the University of California Cooperative Extension's *Riparian Health Assessment for Rangelands (RHAR)*. Published protocols and training opportunities are available for each method, and you should make sure to be familiar with the protocols and properly trained before you undertake these assessments.

The reason for using two assessments is that together they enable you to capture more information regarding the riparian system. There is some overlap between the two assessments, but when you use both you get a comprehensive picture.

It is important to note that not all streams have the same habitat potential. Ward et al. (2001) found that stream morphology affects the streams' habitat potential. For this reason, you should make comparisons only within the same morphology classification. Measurements to determine the *Rosgen classification* (a stream morphology classification system) (Rosgen 1996) should be recorded on the Riparian Grazing Case Study Data Sheet included in this packet.



Figure 3. Team in the field measuring channel morphology cross-sections. To provide an accurate representation, measurements should be taken at every break in slope or every few feet.

PHYSICAL PARAMETERS

The Riparian Grazing Case Study Data Sheet outlines the physical parameters you will have to observe and record. Some equipment is necessary for completion of this data sheet, but if you work with NRCS, RCD, or UCCE offices, this should not be a problem. To begin, you will measure the channel morphology cross-section at both the downstream and upstream transects (transects 1 and 6). Please refer to MacDonald et al. (1991) for the detailed description that begins on page 109. The equipment you will need consists of a stadia rod and scope. Take a reading at every break in slope or every 2 feet (Figure 3). Input the raw data into a computer spreadsheet program and generate a graphical representation of the stream cross-section (see Figures 4 and 5).




Figure 4. Measurements taken in the field can be converted to a graphical representation of the channel morphology. Labeling the banks and thalweg (the deepest part of the channel) helps keep the graph in perspective.




Figure 5. Actual site described graphically in Figure 4. The tape can be seen stretched across the stream, and left and right banks as well as thalweg are highlighted for reference.



Figure 6. Spherical densimeters are used to measure canopy cover.

Take canopy readings along transects, using a densimeter (Figure 6). Again, for specialized equipment and training in its use please contact a local NRCS, RCD, or UCCE office. The densimeter readings will indicate whether canopy cover is increasing.

You also want to document current air and water temperatures, and it is best if you take your readings in the same spot each time. Just select an arbitrary point along one transect and record the location on the data sheet.

HABITAT PARAMETERS

The habitat parameters include calculating the total linear feet of pools (water is deeper and slower moving), riffles (faster and shallower), and runs (sections where water depth and velocity remain more even) (Figure 7). This provides information on the three basic habitat features that are available to fish. In addition, you need to examine specific habitat features. A complete description of all of the parameters under the *Fish Shelter Ratings* section can be found in Flosi et al. (1998).

In determining the percent substrate exposed, you must carefully examine habitat substrates such as boulders, cobbles, woody debris, and the like. This information will vary from year to year with different flow regimes, artificial and natural, but it is important in determining how much habitat is potentially available to fish and macroinvertebrates.



Figure 7. A stretch of stream can contain riffles, runs, and pools, all important features for various habitats. Highlighted in this picture are examples of all three.

Collect specific information regarding three of the riffles in the reach. Consult the protocol for macroinvertebrate collections published by the California Department of Fish and Game (1999) for details. Even though you will not actually sample the macroinvertebrates, the information you collect can provide insight on potential habitat and should certainly be recorded. Length of the riffle as well as average width, depth, and velocity can all easily be recorded with the help of a tape measure, a stopwatch, and a float, such as an orange or a twig. Substrate complexity and embeddedness are examined for each riffle. Using RHAR, *substrate complexity* refers to question 5,

Macroinvertebrate Habitat, and *embeddedness* refers to the High Gradient form, question 9. You will also estimate the percentage of each substrate's size and the degree of its consolidation for each riffle. Finally, you will use a clinometer to determine the gradient of the riffle.

MANAGEMENT SURVEY

Last of all, you will complete a management questionnaire. The Riparian Grazing Case Study Management Survey (Appendix C1) will help you as the manager outline current (*new management*) and historic management (*previous management*) as well as the watershed's characteristics, your goals for the riparian area, and your monitoring practices. The survey should be completed in detail since it will provide a road map of what management practices have been implemented. When you know what management practices are implemented, you have a better idea of what practices may improve a riparian area. Without this information, you will have a hard time comparing manage-

ment changes over time. Complete a new survey each time you change your management methods and you will build up a detailed, useful history.

WHEN SHOULD I REVISIT THE SITE?

You should revisit the case study site on a regular basis, though you will not have to collect data every year. You can expect to repeat the assessments every couple of years, when you implement management changes, or when you notice drastic changes during regular visits to the area.

CONCLUSIONS

By standardizing the data that you collect when you modify riparian grazing management, you will be able to compare various management systems and share ideas with other managers on what management practices have been successful and which have not. This kind of shared experience is one of the best learning opportunities available to land managers. For this reason, it is important that you take the necessary time and care when you gather your data. If you collect good data at the beginning, you can put it to good use for years to come.

REFERENCES

- California Department of Fish and Game Aquatic Bioassessment Laboratory. 1999. California stream bioassessment procedure: Habitat and biological assessment for citizen monitors. Rancho Cordova, CA: Water Pollution Control Laboratory.
<http://www.dfg.ca.gov/cabw/cabw/csbg.html>
- Flosi, G., S. Downie, J. Hopelain, M. Bird, R. Coey, and B. Collins. 1998. California salmonid stream habitat restoration manual. Third edition. Sacramento: California Department of Fish and Game.
<http://www.dfg.ca.gov/habitats/index.html>
- MacDonald, L. H., A. W. Smart, and R. C. Wissmar. 1991. Monitoring guidelines to evaluate effects of forestry activities on streams in the Pacific Northwest and Alaska. Washington, DC: U.S. Environmental Protection Agency.
- Natural Resources Conservation Service. 1998. Stream visual assessment protocol. NWCC-TN-99-1. Portland, OR: National Water and Climate Center.
<http://www.wcc.nrcs.usda.gov/water/quality/common/nrcsdocs.html>
- Prichard, D., J. Anderson, C. Correll, J. Fogg, K. Gebhardt, R. Krapf, S. Leonard, B. Mitchell, and J. Staats. 1998. Riparian area management TR 1737-15: A user guide to assessing proper functioning condition and the supporting science for lotic areas. Denver: U.S. Bureau of Land Management, National Business Center, BC-650B, P.O. Box 25047, Denver, CO 80225-0047.
- Rosgen, D. 1996. Applied river morphology. Wildland Hydrology.
- Ward, T. A., K. W. Tate, and E. R. Atwill. 2001. A comparison of three riparian resource assessments. Society for Range Management, 54th Annual Conference.
- Ward, T. A., K. W. Tate, and E. R. Atwill. 2003. Visual assessment of riparian health. Oakland: University of California Division of Agriculture and Natural Resources Publication 8089.
<http://anrcatalog.ucdavis.edu/pdf/8089LR.pdf>
- Winward, A. H. 2000. Monitoring the vegetation resources in riparian areas. Washington, DC: USDA Forest Service.

Appendix A
Riparian Grazing Case Study Monitoring Tools

Assessment	Short- or long-term trend	Parameter quantified
BLM Proper Functioning Condition	Long-term	Hydrologic function
UCCE Riparian Health Assessment for Rangelands	Long-term	Trout and macroinvertebrate habitat and hydrologic function
Greenline: Vegetation Cross-section Composition	Short- and long-term	Width of the riparian area
Greenline: Greenline Composition	Short- and long-term	Change in greenline vegetation
Greenline: Woody Species Regeneration	Short- and long-term	Change in woody species along the greenline
Channel Morphology Cross-section	Long-term	Change in width and depth of the channel
Densiometer	Long-term	Amount of canopy
Habitat Types	Short- and long-term	Three basic habitat types for fish
Physical Parameters	Short- and long-term	Variety of physical parameters

Appendix B

Check Sheet for Establishing Case Studies

I. Ahead of time

- a. Review protocols
- b. Receive training if necessary
- c. Gather required equipment
 - i. Stadia rod
 - ii. Hand lens
 - iii. Tape (300-ft if possible)
 - iv. Densiometer
 - v. Stakes, sledgehammer, and paint
 - vi. Compass
 - vii. Clinometer
 - viii. Make copies of necessary forms
 1. Six Vegetation Cross-section sheets
 2. One Greenline form
 3. One Woody Species Regeneration form
 4. One RHAR form
 5. One PFC form
 6. One Riparian Case Study Data Sheet
 7. One Riparian Grazing Case Study Management Survey

II. At site

- a. Select representative section within one Rosgen type
- b. Establish transects (Figure 1)
 - i. Record distance and bearing from benchmark
- c. Begin Greenline (three parts)
 - i. Vegetation cross-section transects
 1. Record canopy reading on densiometer at mid-channel for each transect
 - ii. Greenline composition transect
 - iii. Woody species regeneration belt transect
- d. Complete visual assessments (RHAR and PFC)
- e. Complete channel morphology cross-sections (Transects 1 and 6)
- f. Complete Riparian Grazing Case Study Data Sheet
 - i. Air and water temperature
 - ii. Stream morphology data (widths and depths)
 - iii. Feet of riffles, pools, and runs
 - iv. Fish shelter ratings
 - v. Riffle data
- g. Complete Management Survey

Appendix C1

Riparian Grazing Case Study Management Survey page 1

General Information

Ranch: _____
Name: _____
Address: _____
City, State, ZIP: _____
Phone number: _____
E-mail: _____

Ownership:
Private Private lease
U.S. Forest Service BLM
Other public
How long under current ownership?

If public-owned, is there regular communication with USFS or BLM Range Con ?
Yes No

Type of operation:
Cow-calf Stocker Sheep
Farming Horses
Total size and number of pastures:

Watershed Characteristics

Upstream watershed land uses:
Urban Logging
Ranching Farming
Wildlands Recreation
Roads Non-urban residential

Predominant ownership of watershed:
Private U.S. Forest Service
BLM Public

Past land disturbances in the watershed:
Mining Floods Fire
Logging Landslides

Management Unit of Concern

Name: _____
County: _____
Ownership:
Private Private lease
U.S. Forest Service BLM
Other public
How long under current ownership?

If public-owned, are there standards in place?
Yes No

What are the standards?
Utilization % Stubble height inches
Browse % Trampling %
RDM lbs/acre

Who monitors them?
Range Con Rancher

Size and number of pastures in unit:
Acres: _____
Number of pastures: _____

How many pastures contain a section of creek?

Are there any written plans for the unit?
Ranch plan Water quality plan
Economic plan EQIP
AOI EA/EIS; IS/EIR
Conservation agreement
Land use plan Other

Goals for riparian pasture:
Increase/maintain production
Increase/maintain profit
Maintain/improve water quality
Aesthetics
Sustainability
Increase biodiversity
Decrease weeds
Improve/maintain fishery

Have you created a separate riparian pasture specifically to obtain achieve your goals?
Yes No

If yes, how long did you allow the new pasture to rest before grazing was reintroduced?
One season One year
Two years Three years
Four or more years

Are temporary exclosures utilized to meet your goals in the riparian area?
Yes No

Riparian concerns that you have:
Fish habitat Wildlife habitat
Waterfowl habitat Water quality

Riparian Grazing Case Study Management Survey page 2

Biomass production TMDL

Endangered Species Act

Use of pasture:

Holding area Calving

Watering site Grazing

Gathering Bedding

Exclosure

Indicators used to move livestock in and out of riparian area (unit of concern):

- Dormant season of key plants
- Invasion of undesirable plants/Shading of desirables
- Bank soil moisture
- Presence and/or life cycle of key wildlife species?
- Browse on key woody vegetation
- Accumulation of liter layer
- RDM level
- Likelihood of floods/spring runoff
- Utilization of herbaceous vegetation
- Time of year (calendar dates)
- Rest period of other pastures

Current Management, Costs (days of labor/year), and Possible Cost Sharing, (for the particular pasture, not the entire ranch)

Type of operation and length of time under current operation.

- Cow-calf Stocker Sheep
- Farming Horses

Breed/type of animal: _____

Number of animals (range and average): _____

Season of use:

- Spring Summer
- Fall Winter

Average in and out dates, or time between rotations: _____

Grazing system: _____

Livestock distribution

- Herding Drift fence
- Trails Temporary exclosures
- Off-site Feed or Salt/minerals

If you use off-site feeding and/or salt/minerals:

How far is the off-site feed/salt/minerals from the stream? (closest 1/2 mile is fine) _____

Do you observe evidence of livestock using off-site feed/salt/minerals?

- Yes No

In your opinion/observation has the off-site feed/salt/minerals reduced time livestock spend in the riparian area?

- Yes No

Is off-site water available:

- Yes No

If yes: Natural Human-made

Type of human-made:

- Pipeline Troughs
- Tanks Well
- Pond

How far is the off-site water from the stream? (closest 1/2 mile is fine)

Do you observe evidence of livestock using off-site water?

- Yes No

In your opinion/observation has the off-site water reduced time livestock spend in the riparian area?

- Yes No

Brush Management

- Fire Chemical Mechanical

Are you performing brush management practices to obtain/achieve your riparian goals?

- Yes No

Riparian Grazing Case Study Management Survey page 3

Road Management

- Maintenance Construction
- Culverts

Are you performing road management practices to obtain/ achieve your riparian goals?

- Yes No

Fencing

Type of fencing used:

- Barbed wire Electric, 5-strand
- Electric, 3-strand Electric, 2-strand
- Electric, 1-strand Temp. electric

- Range seeding

Stream crossings (interim):

- For livestock
- For roads (equipment, truck)
- If for livestock, are they hardened?
- Yes No

How often are they utilized?

Have they reduced damage to the stream banks in your opinion?

- Yes No

If for roads, are they hardened?

- Yes No

How often are they used?

Are they County? Private?

- Prescribed burning for forage improvement
- Irrigation water management
- Pasture clipping
- Sediment basins
- Grazingland mechanical treatments (renovating, contour furrowing, pitting)
- Length of time under current management?

Restoration Efforts

Has there been any restoration in the unit?

- Yes No

If so, what was the objective?

- Decrease erosion
- Capture sedimentation
- Improve habitat
- Sustainability of the system

What restoration practices were utilized?

- Stream corridor improvement
- Bank protection
- Structural (such as rock riprap)
- Bioengineering(either solely vegetation such as willows, or a combination of vegetation and structural)
- Stream channel stabilization
- Grade stabilization
- Riparian planting for wildlife habitat
- Wildlife habitat in the upland
- Critical area planting for erosion
- Landslide treatments
- Do you purposely cull animals that "hug the stream" ("riparian huggers")?
- Does anyone stock fish?

Historic Management and Costs (for the particular area)

Type of operation and length of time under historic operation.

- Cow-calf Stocker Sheep
- Farming Horses

Breed/type of animal: _____

Number of animals (range and average):

Season of use:

- Spring Summer
- Fall Winter

Average in and out dates, or time between rotations:

Grazing system description:

Riparian Grazing Case Study Management Survey page 4

Livestock distribution

- Herding
- Trails
- Off-site Feed or Salt/minerals
- Drift fence
- Temporary enclosures

If you used off-site feeding and/or salt/minerals, How far was the off-site feed/salt/minerals from the stream? (closest 1/2 mile is fine) _____

Did you observe evidence of livestock using off-site feed/salt/minerals?
 Yes No

In your opinion/observation did the off-site feed/salt/minerals reduced time livestock spend in the riparian area?
 Yes No

Was off-site water available:
 Yes No

If yes: Natural Human-made
 Type of human-made:
 Pipeline Troughs
 Tanks Well
 Pond

How far was the off-site water from the stream? (closest 1/2 mile is fine)

Did you observe evidence of livestock using off-site water?
 Yes No

In your opinion/observation did the off-site water reduced time livestock spend in the riparian area?
 Yes No

Brush management (314)
 Fire Chemical Mechanical

Did you performing brush management practices to obtain/achieve your riparian goals?
 Yes No

Road management
 Maintenance Construction
 Culverts

Did you performing road management practices to obtain/achieve your riparian goals?
 Yes No

Fencing (382)
 Type of fencing used:
 Barbed wire Electric, 5-strand
 Electric, 3-strand Electric, 2-strand
 Electric, 1-strand Temp. electric

Range Seeding

Stream crossings (interim):
 For livestock
 For roads (equipment, truck)
 If for livestock, were they hardened?
 Yes No

How often were they utilized?

Did they reduced damage to the stream banks in your opinion?
 Yes No

If for roads, were they hardened?
 Yes No

How often are they used? _____
 Are they County? Private?

- Prescribed burning for forage improvement
- Irrigation water management
- Pasture clipping
- Sediment basins
- Grazingland mechanical treatments (renovating, contour furrowing, pitting)
- Length of time under historic management?

Restoration Efforts

Was there any historic restoration in the unit?
 Yes No

If so, what was the objective?
 Decrease erosion
 Capture sedimentation
 Improve habitat

Riparian Grazing Case Study Management Survey page 5

- Sustainability of the system
- What restoration practices were utilized:
- Stream corridor improvement
- Bank protection
- Structural (such as rock riprap)
- Bioengineering(either solely vegetation such as willows, or a combination of vegetation and structural)
- Stream channel stabilization
- Grade stabilization
- Riparian planting for wildlife habitat
- Wildlife habitat in the upland
- Critical area planting for erosion
- Landslide treatments
- Did you purposely cull animals that “hug the stream” (Riparian Huggers)?
- Did anyone stock fish? _____

Current Monitoring

Types of monitoring, number of points and how often:

	Frequency (per yr)	Location
<input type="checkbox"/> Visual:	_____	_____
<input type="checkbox"/> Photo:	_____	_____
<input type="checkbox"/> Stream temp:	_____	_____
<input type="checkbox"/> Sediment	_____	_____
<input type="checkbox"/> Nutrient	_____	
<input type="checkbox"/> Habitat:	_____	
<input type="checkbox"/> Pathogens:	_____	
<input type="checkbox"/> Wildlife:	_____	

Objectives of monitoring:

- Establish base lines
- Document management over time
- Monitor wildlife/fisheries habitat
- Monitor vegetation: weeds and desirable grasses
- Protect ranching interests against environmental concerns

How are monitoring data used?

- To make management decisions
- Stored for future use
- Shared with agencies (Regional Board, NRCS, UCCE, RCD, FS, BLM, F&G, etc.)

Appendix C2
Filled-in Example of Appendix C1

Riparian Grazing Case Study Management Survey page 1

General Information

Ranch: UCC Example Site
Name: Agronomy, Range Science
Address: One Shields Ave
City, State, ZIP: Davis, CA 95616
Phone number: (530) 752-4031
E-mail: _____

Ownership:
 Private Private lease
 U.S. Forest Service BLM
 Other public

How long under current ownership?
98 years

If public-owned, is there regular communication with USFS or BLM Range Con ?
 Yes No

Type of operation:
 Cow-calf Stocker Sheep
 Farming Horses

Total size and number of pastures:
3425 acres 15 pastures

Watershed Characteristics

Upstream watershed land uses:
 Urban Logging
 Ranching Farming
 Wildlands Recreation
 Roads Non-urban residential

Predominant ownership of watershed:
 Private U.S. Forest Service
 BLM Public

Past land disturbances in the watershed:
 Mining Floods Fire
 Logging Landslides

Management Unit of Concern

Name: UCC Example Creek
County: Yolo
Ownership:
 Private Private lease
 U.S. Forest Service BLM
 Other public
How long under current ownership?
98 years

If public-owned, are there standards in place?
 Yes No

What are the standards?
 Utilization ____% Stubble height ____inches
 Browse ____% Trampling ____%
 RDM ____lbs/acre

Who monitors them?
 Range Con Rancher

Size and number of pastures in unit:
Acres: 248
Number of pastures: 1
How many pastures contain a section of creek?
1

Are there any written plans for the unit?
 Ranch plan Water quality plan
 Economic plan EQIP
 AOI EAVEIS, IS/EIR
 Conservation agreement
 Land use plan Other

Goals for riparian pasture:
 Increase/maintain production
 Increase/maintain profit
 Maintain/improve water quality
 Aesthetics
 Sustainability
 Increase biodiversity
 Decrease weeds
 Improve/maintain fishery

Have you created a separate riparian pasture specifically to obtain achieve your goals?
 Yes No

If yes, how long did you allow the new pasture to rest before grazing was reintroduced?
 One season One year
 Two years Three years
 Four or more years

Are temporary exclosures utilized to meet your goals in the riparian area?
 Yes No

Riparian concerns that you have:
 Fish habitat Wildlife habitat
 Waterfowl habitat Water quality

Appendix C2
Filled in Example of Appendix C1 (continued)

Riparian Grazing Case Study Management Survey page 2

Biomass production TMDL

Endangered Species Act

Use of pasture:

Holding area Calving

Watering site Grazing

Gathering Bedding

Exclosure

Indicators used to move livestock in and out of riparian area (unit of concern):

Dormant season of key plants

Invasion of undesirable plants/Shading of desirables

Bank soil moisture

Presence and/or life cycle of key wildlife species?

Browse on key woody vegetation

Accumulation of litter layer

RDM level

Likelihood of floods/spring runoff

Utilization of herbaceous vegetation

Time of year (calendar dates)

Rest period of other pastures

Current Management, Costs (days of labor/year), and Possible Cost Sharing, (for the particular pasture, not the entire ranch)

Type of operation and length of time under current operation.

Cow-calf Stocker Sheep

Farming Horses

for 98 years

Breed/type of animal: Angus

Number of animals (range and average):

50-100 normally 72

Season of use:

Spring Summer

Fall Winter

Average in and out dates, or time between rotations:

10/15 → 11/1

Grazing system:

Used in fall for two weeks every year. Cattle excluded rest

of the year.

Livestock distribution

Herding Drift fence

Trails Temporary exclosures

Off-site Feed or Salt/minerals 1 day/yr

If you use off-site feeding and/or salt/minerals:

How far is the off-site feed/salt/minerals from the stream? (closest 1/2 mile is fine) 3/4 mile

Do you observe evidence of livestock using off-site feed/salt/minerals?

Yes No

In your opinion/observation has the off-site feed/salt/minerals reduced time livestock spend in the riparian area?

Yes No

Is off-site water available:

Yes No

If yes: Natural Human-made

Type of human-made:

Pipeline Troughs

Tanks Well

Pond

How far is the off-site water from the stream? (closest 1/2 mile is fine)

1/2 mile

Do you observe evidence of livestock using off-site water?

Yes No

In your opinion/observation has the off-site water reduced time livestock spend in the riparian area?

Yes No

Brush Management

Fire Chemical Mechanical

Are you performing brush management practices to obtain/achieve your riparian goals?

Yes No

Appendix C2
 Filled in Example of Appendix C1 (continued)

Riparian Grazing Case Study Management Survey page 5

Sustainability of the system

What restoration practices were utilized:

- Stream corridor improvement
- Bank protection
- Structural (such as rock riprap)
- Bioengineering (either solely vegetation such as willows, or a combination of vegetation and structural)
- Stream channel stabilization
- Grade stabilization
- Riparian planting for wildlife habitat
- Wildlife habitat in the upland
- Critical area planting for erosion
- Landslide treatments
- Did you purposely cull animals that "hug the stream" (Riparian Huggers)?
- Did anyone stock fish? _____

Current Monitoring

Types of monitoring, number of points and how often:

	Frequency (per yr)	Location
<input checked="" type="checkbox"/> Visual:	<i>daily when cattle present, plus 4 times throughout year.</i>	
<input checked="" type="checkbox"/> Photo:	<u>2</u>	<u>4 spots</u>
<input type="checkbox"/> Stream temp:	_____	_____
<input type="checkbox"/> Sediment	_____	_____
<input type="checkbox"/> Nutrient	_____	_____
<input checked="" type="checkbox"/> Habitat:	<i>once every other year</i>	
<input type="checkbox"/> Pathogens:	_____	_____
<input checked="" type="checkbox"/> Wildlife:	<i>visual counts</i>	

Objectives of monitoring:

- Establish base lines
- Document management over time
- Monitor wildlife/fisheries habitat
- Monitor vegetation: weeds and desirable grasses
- Protect ranching interests against environmental concerns

How are monitoring data used?

- To make management decisions
- Stored for future use
- Shared with agencies (Regional Board, NRCS, UCCE, RCD, FS, BLM, F&G, etc.)

Appendix C3 (continued)
Riparian Grazing Case Study Data Sheet page 2

Transect Locations: (Lat., Long., distance and bearing from Bench Mark, etc.)

Tran. 1: _____

Tran. 2: _____

Tran. 3: _____

Tran. 4: _____

Tran. 5: _____

Tran. 6: _____

	Riffle 1	Riffle 2	Riffle 3	Fish Shelter Ratings											
Riffle Length	_____	_____	_____	% undercut bank	<table border="1"> <tr><td> </td></tr> </table>										
Avg. Riffle Length	_____	_____	_____	% swd											
Avg. Riffle Depth	_____	_____	_____	% lwd											
Riffle Velocity	_____	_____	_____	% root mass											
Substrate Complexity	_____	_____	_____	% terr. veg											
Embeddedness	_____	_____	_____	% aqua. veg											
Substrate Composition	% Fines _____ % Gravel _____ % Cobble _____ % Boulder _____ % Bedrock _____	_____ _____ _____ _____ _____	_____ _____ _____ _____ _____	% boulder curtain											
				% boulder											
				% bedrock ledge											
				% Exposed Substrate											
				<table border="1"> <tr><td> </td></tr> </table>											
Substrate Consolidation	_____	_____	_____												
% Gradient	_____	_____	_____												

Appendix C4 (page 2)
 Filled in Example of Appendix C3

Riparian Grazing Case Study Data Sheet

Transect Locations: (Lat., Long., distance and bearing from Bench Mark, etc.)

- Tran. 1: N41° 06' 01.8" W120° 22' 11.3" 10ft from rock outcrop bearing 210°
- Tran. 2: N41° 05' 02.2" W120° 22' 10.1" 23.5ft from bench mark bearing 198°
- Tran. 3: N41° 05' 04.3" W120° 22' 8.8" 39.4ft from bench mark bearing 183°
- Tran. 4: N41° 04' 08.4" W120° 22' 07.9" 56.2ft from bench mark bearing 179°
- Tran. 5: N41° 04' 07.9" W120° 22' 06.5" 92.4ft from bench mark bearing 163°
- Tran. 6: N41° 04' 05.5" W120° 22' 05.3" 121.6ft from bench mark bearing 167°

	Riffle 1	Riffle 2	Riffle 3
Riffle Length	<u>17.2'</u>	<u>15'</u>	<u>8'</u>
Avg. Riffle Length	<u>5'</u>	<u>7'</u>	<u>8'</u>
Avg. Riffle Depth	<u>0.33'</u>	<u>0.33'</u>	<u>0.5'</u>
Riffle Velocity	<u>2.8 ft/sec</u>	<u>2.1 ft/sec</u>	<u>1.5 ft/sec</u>
Substrate Complexity	<u>16</u>	<u>16</u>	<u>17</u>
Embeddedness	<u>20%</u>	<u>40%</u>	<u>10%</u>
Substrate Composition			
% Fines	<u>15</u>	<u>25</u>	<u>10</u>
% Gravel	<u>30</u>	<u>35</u>	<u>55</u>
% Cobble	<u>55</u>	<u>40</u>	<u>15</u>
% Boulder	<u>-</u>	<u>-</u>	<u>20</u>
% Bedrock	<u>-</u>	<u>-</u>	<u>-</u>
Substrate Consolidation	<u>loose</u>	<u>moderate</u>	<u>loose</u>
% Gradient	<u>2%</u>	<u>2%</u>	<u>2%</u>

Fish Shelter Ratings	
% undercut bank	<u>0</u>
% swd	<u>5</u>
% hwd	<u>0</u>
% root mass	<u>0</u>
% terr. veg	<u>50% of bank covered</u>
% aqua. veg	<u>50%</u>
% boulder curtain	<u>0</u>
% boulder	<u>5</u>
% bedrock ledge	<u>0</u>
% Exposed Substrate	<u>5%</u>

Appendix C5
Riparian Greenline Transect Data Form

RIPARIAN GREENLINE TRANSECT DATA

Forest / District _____ / _____ Date 9/2/01
 Drainage VCE Example Creek
 Examiners David, Don, Theresa Photo No's _____
 Complex _____
 Location _____
 Transect No. _____ Feet/Step 3/1

Community Type	STEPS (Left)				STEPS (Right)					TOTAL STEPS	% COMP.			
	bare bank / Rabbit Brush	Per grass	Sedge											
bare bank / Rabbit Brush / Per grass / Sedge	35	13	1	14				3	1	5	4	5	81	26.8
Rush / Per grass	8	7						5					20	6.6
Per grass / Sedge / Rush	14	52	9					32	6	13	5	70	201	66.6
Grand Total											302			

BARS WITHIN TRANSECT (Optional)

	STEPS	FEET
GRAVEL		
SAND		
SILT/CLAY		

Total Steps ea. CT

Grand Total = Composition

Appendix C6 (page 1)
 Low-Gradient Riparian Health Assessment for Rangelands Form

Low Gradient Riparian Health Assessment for Rangelands

Site: UCCE Example Creek Date: 9/2/01

1. Channel Condition	Natural channel, no evidence of down cutting.	Evidence of past channelization or down cutting, but significant recovery. Adequate access to the flood plain.	Channelization or down cutting extensive. Flood plain is restricted.	Channel actively down cutting or widening. Flood plain access prevented.
Score:	12 11 10	9 8 7	6 <u>5</u> 4	3 2 1
2. Access to Flood Plain	Flooding every 1 1/2 to 2 years -- not incised	Flooding every 3-5 years -- limited incision	Flooding every 6-10 years -- deeply incised	No flooding. Deeply incised.
Score:	12 11 10	9 8 <u>7</u>	6 5 4	3 2 1
3. Bank stability (each bank separately -Looking downstream, Left bank - Right bank)	Banks stable. Outside bends protected by roots.	Moderately stable. Infrequent, small areas of erosion, mostly healed over.	Moderately unstable. Outside bends actively eroding; banks high; high potential erosion.	Unstable. Actively eroding.
Left Bank Score:	6 5.5 5	<u>4.5</u> 4 3.5	3 2.5 2	1.5 1 0.5
Right Bank Score:	6 5.5 5	4.5 <u>4</u> 3.5	3 2.5 2	1.5 1 0.5
4a. Riparian Zone-Perennial Creek (score each bank separately)	Natural vegetation extends at least two active channel widths (ex. sedges, rushes, willows, alders, aspen, cottonwoods, sycamores). Score higher if point bars are re-vegetating and all age classes of woody species present (seedling, young, mature, old)	Natural vegetation extends one active channel width. Or Covers flood plain.	Natural vegetation extends 1/2 active width. Or Filtering function moderately compromised.	Natural vegetation extends less than 1/2 of active width. Or Lack of regeneration. Or Filtering function severely compromised.
Left Bank Score:	6 5.5 5	4.5 4 <u>3.5</u>	3 2.5 2	1.5 1 0.5
Right Bank Score:	6 5.5 5	4.5 4 3.5	<u>3</u> 2.5 2	1.5 1 0.5
4a. Riparian Zone-Intermittent Creek (score each bank separately)	Natural vegetation extends two active channel widths (ex. oaks, buckeyes, alders, cottonwoods, annual grasses, some sedges and rushes). Score higher if point bars are re-vegetating and all age classes of woody species present (seedling, young, mature, old)	Natural vegetation extends one active channel width. Bare spots common. Or Covers flood plain.	Natural vegetation extends 1/2 of active width. Bare spots common. Or Filtering function moderately compromised.	Natural vegetation extends less than 1/2 of active width. Or Lack of regeneration. Or Filtering function severely compromised.
Left Bank Score:	6 5.5 5	4.5 4 3.5	3 2.5 2	1.5 1 0.5
Right Bank Score:	6 5.5 5	4.5 4 3.5	3 2.5 2	1.5 1 0.5

Appendix C6 (page 2)
 Low-Gradient Riparian Health Assessment for Rangelands Form

5. Macroinvertebrate Habitat	Greater than 5 habitat types. Score higher if good diversity.	3-4 types	1-2 types	0-1 type
Cover types:	<u>Boulders, cobbles, coarse gravel</u> , leaf packs, <u>fine woody debris</u> , submerged logs, overhanging vegetation, <u>macrophytes</u> (aquatic vegetation)			
Score:	12 11 <u>10</u>	9 8 7	6 5 4	3 2 1
6. Macroinvertebrates Observed	Class I dominate. Score higher if good diversity and number.	Class II dominate.	Class III dominate.	No macroinvertebrates present.
Score:	12 <u>11</u> 10	9 8 7	6 5 4	3 2 1
7. Fish Habitat (if applicable)	Greater than 7 habitat type. Score higher if good diversity.	6-4 habitats present.	3-2 habitats present.	1-0 habitat present.
Cover types:	Logs/large woody debris, deep pools, overhanging vegetation, <u>riffles</u> , <u>boulders/cobbles</u> , thick root mats, isolated/backwater pools, <u>dense macrophyte beds</u> , undercut banks			
Score:	12 11 10	9 8 7	6 <u>5</u> 4	3 2 1
8. Pool Variability	Even mix large-shallow, large-deep, small-shallow, small-deep pools present.	Majority of pools large-deep.	Shallow pools more prevalent than deep pools.	Majority of pools small-shallow or pools absent.
Score:	12 11 10	9 8 7	6 5 4	<u>3</u> 2 1
9. Pool Substrate	Mix of substrate (gravel, firm sand, etc.). Roots, submerged vegetation common.	Mix of soft sand, mud, and clay. Some submerged vegetation.	All mud, clay, or sand. Little to no root mats or submerged vegetation.	Hard-pan clay or bedrock. No roots or submerged vegetation.
Score:	12 11 10	<u>9</u> 8 7	6 5 4	3 2 1
10. Channel Flow	Water reaches base of both lower banks, minimal substrate exposed.	Water fills >75% of the channel, <25% of substrate exposed.	Water fills 25-75% of the channel, riffle substrate mostly exposed.	Very little water in channel and mostly present in standing pools.
Score:	12 <u>11</u> 10	9 8 7	6 5 4	3 2 1

Photopoint Monitoring

	Location Description	Compass Heading	Landmarks
Witness Point			
Photopoint 1			
Photopoint 2			
Photopoint 3			

Date /time	Photographer	Photo-point #	Camera/lens/ film speed	Roll #/ frame #	Observations

total score: 7.75

Appendix C7 (page 1)
PFC Standard Checklist Form

Standard Checklist

Name of Riparian-Wetland Area: VCC E Example Creek

Date: 9/2/01 Segment/Reach ID: _____

Miles: _____ Acres: _____

ID Team Observers: _____

Yes	No	N/A	HYDROLOGIC
✓			1) Active floodplain inundated in "relatively frequent" events (1-3 years) <i>current floodplain is.</i>
		✓	2) Active/stable beaver dams <i>beaver activity but no dams.</i>
✓			3) Sinuosity, width/depth ratio, and gradient are in balance with the landscape setting (i.e., landform, geology, and bioclimatic region)
✓			4) Riparian area is widening or has achieved potential extent <i>is starting to widen.</i>
✓			5) Upland watershed is not contributing to riparian degradation

Yes	No	N/A	VEGETATION
	✓		6) Diverse age-class distribution (recruitment for maintenance/recovery) <i>no mature/old class</i>
✓			7) Diverse composition of vegetation (for maintenance/recovery)
✓			8) Species present indicate maintenance of riparian soil moisture characteristics
✓			9) Streambank vegetation is made up of those plants or plant communities that have root masses capable of withstanding high streamflow events
	X		10) Riparian plants exhibit high vigor <i>beaver activity on the willows</i>
	✓		11) Adequate vegetative cover present to protect banks and dissipate energy during high flows
✓			12) Plant communities in the riparian area are an adequate source of coarse and/or large woody debris

Appendix C8 (page 1)

Six Cross Section Composition Forms and One Cross Section Summary Sheet

**CROSS SECTION COMPOSITION
(Transect Data)**

Forest / District _____ / _____ Date 9/2/01

Drainage UCCE Example Creek

Examiners David, Theresa, Don Photo No's _____

Complex _____

Location _____

Transect No. 1 Feet/Step 31

Community Type	NUMBER STEPS										TOTAL STEPS	FEET Optional	
	1	2	3	4	5	6	7	8	9	10			
per grass / Rabbit Brush / Sedge mix	7	5										12	36
bare ground / per grass / Rabbit brush	5											5	15
per grass / Sedge / Rush / Forb	4											4	12
Creek	2											2	6
per grass / Rush / Forb	1											1	3
Bare ground	2											2	6

ESTIMATED AVERAGE HT.	Sprout	Young	Mature	Decadent	Dead

LINE INTERCEPT CANOPY OF WOODY SPECIES (optional) _____

TOTAL FEET OF RIPARIAN (optional) _____

Appendix C8 (page2)
Six Cross Section Composition Forms and One Cross Section Summary Sheet

**CROSS SECTION COMPOSITION
(Transect Data)**

Forest / District _____ / _____ Date 9/2/01
 Drainage ULCE Example Creek
 Examiners David, Don, Theresa Photo No's _____
 Complex _____
 Location _____
 Transect No. 2 Feet/Step 3/1

Community Type	NUMBER STEPS										TOTAL STEPS	FEET Optional
	1	2	3	4	5	6	7	8	9	10		
Barc ground / Rabbit brush	1	2	3	4	5	6	7	8	9	10	12	36
Per grass / Rabbit brush	1	2	3	4	5	6	7	8	9	10	13	39
Per grass / Rush / Forb	1	2	3	4	5	6	7	8	9	10	3	9
Creek	1	2	3	4	5	6	7	8	9	10	2	6
barc ground	1	2	3	4	5	6	7	8	9	10	3	9

ESTIMATED AVERAGE HT.	Sprout	Young	Mature	Decadent	Dead

LINE INTERCEPT CANOPY OF WOODY SPECIES (optional) _____

TOTAL FEET OF RIPARIAN (optional) _____

Appendix C8 (page3)

Six Cross Section Composition Forms and One Cross Section Summary Sheet

**CROSS SECTION COMPOSITION
(Transect Data)**

Forest / District _____ / _____ Date 9/2/01

Drainage ULCE Example Creek

Examiners David, Don, Theresa Photo No's _____

Complex _____

Location _____

Transect No. 3 Feet/Step 3/1

Community Type	NUMBER STEPS										TOTAL STEPS	FEET Optional
	1	2	3	4	5	6	7	8	9	10		
Bare / Per / Rabbit ground / grass / Brush	8	6									14	42
Rush / Sedge / Per grass	4	1	3								8	24
Willow	1	1									2	6
Creek	2										2	6
bare ground	3										3	9

ESTIMATED AVERAGE HT.	Sprout	Young	Mature	Decadent	Dead
		4'			

LINE INTERCEPT CANOPY OF WOODY SPECIES (optional) _____

TOTAL FEET OF RIPARIAN (optional) _____

Appendix C8 (page 4)
 Six Cross Section Composition Forms and One Cross Section Summary Sheet

**CROSS SECTION COMPOSITION
 (Transect Data)**

Forest / District _____ / _____ Date 9/2/01
 Drainage UCCE Example Creek
 Examiners David, Don, Theresa Photo No's _____
 Complex _____
 Location _____
 Transect No. 4 Feet/Step 3/1

Community Type	NUMBER STEPS										TOTAL STEPS	FEET Optional
	1	2	3	4	5	6	7	8	9	10		
Bare ground / Per grass / Rabbit Brush	9	4	6								19	57
Rush / Per grass / Bare ground	1										1	3
Creek	3										3	9
Rush	1										1	3
Willow	2										2	6
Per grass / Sedge / Rush	4										4	12

ESTIMATED AVERAGE HT.	Sprout	Young	Mature	Decadent	Dead
		36'			

LINE INTERCEPT CANOPY OF WOODY SPECIES (optional) _____

TOTAL FEET OF RIPARIAN (optional) _____

Appendix C8 (page 5)

Six Cross Section Composition Forms and One Cross Section Summary Sheet

**CROSS SECTION COMPOSITION
(Transect Data)**

Forest / District _____ / _____ Date 9/2/01
 Drainage UCC Example Creek
 Examiners David, Don, Theresa Photo No's _____
 Complex _____
 Location _____
 Transect No. 5 Feet/Step 3/1

Community Type	NUMBER STEPS										TOTAL STEPS	FEET Optional
	1	2	3	4	5	6	7	8	9	10		
Bare / Sage ground / Brush	9	1									13	36
Per grass / Sage / Brush	2	4									6	18
Per grass / Rush	1	1									2	6
Creek	3										3	9
Willow	2										2	6
Sedge / Per grass / Rush	3										3	9

ESTIMATED AVERAGE HT.	Sprout	Young	Mature	Decadent	Dead
			5'		

LINE INTERCEPT CANOPY OF WOODY SPECIES (optional) _____

TOTAL FEET OF RIPARIAN (optional) _____

Appendix C8 (page 6)
 Six Cross Section Composition Forms and One Cross Section Summary Sheet

**CROSS SECTION COMPOSITION
 (Transect Data)**

Forest / District _____ / _____ Date 9/2/01
 Drainage UCE Example Creek
 Examiners David, Don, Theresa Photo No's _____
 Complex _____
 Location _____
 Transect No. 6 Feet/Step 3/1

Community Type	NUMBER STEPS										TOTAL STEPS	FEET Optional
	1	2	3	4	5	6	7	8	9	10		
<u>Per grass / Sedge / Sage / Brash</u>	<u>5</u>	<u>2</u>									<u>7</u>	<u>21</u>
<u>Bare ground / Sage / Brash</u>	<u>5</u>										<u>5</u>	<u>15</u>
<u>Willow</u>	<u>2</u>										<u>2</u>	<u>6</u>
<u>Creek</u>	<u>2</u>										<u>2</u>	<u>6</u>
<u>Sedge / Forb / Rush / Per grass</u>	<u>4</u>										<u>4</u>	<u>12</u>
<u>Bareground</u>	<u>2</u>	<u>1</u>	<u>3</u>								<u>6</u>	<u>18</u>
<u>Rosebush</u>	<u>1</u>										<u>1</u>	<u>3</u>

ESTIMATED AVERAGE HT.	Sprout	Young	Mature	Decadent	Dead
			<u>14'</u>		

LINE INTERCEPT CANOPY OF WOODY SPECIES (optional) _____
 TOTAL FEET OF RIPARIAN (optional) _____

Appendix C8 (page 7)
Six Cross Section Composition Forms and One Cross Section Summary Sheet

CROSS SECTION SUMMARY SHEET

Forest/District _____ / _____ Date Compiled 9/2/01

Drainage ULLE Example Creek

Examiners David, Don, Theresa

Complex _____

Transect No's _____

Community Type	T ₁ Steps	T ₂ Steps	T ₃ Steps	T ₄ Steps	T ₅ Steps	T ₆ Steps	PCT COMPOSITION
Per grass / Rabbit / sedge Brush / mix	12						7.5
Bare ground / Per grass / Rabbit Brush	5		14	19			23.8
Per grass / Sedge / Rush / Forb	4					4	5
Bare ground / sage Brush					13	5	11.3
Per grass / Rush / Forb	1	3					2.5
Bareground	2	3	3			6	8.8
Bare ground / Rabbit Brush		12					7.5
Per grass / Rabbit Brush		13					8.1
Rush / Sedge / Per grass			8	4	3		9.4
Willow			2	2	2	2	5
Rush / Per grass / Bare ground				1			0.6
Rush				1			0.6
Per grass / Rush					2		1.2
Per grass / Sage Brush					6		3.8
Rosebush						1	0.6
Per grass / Sedge / Sage Brush						7	4.4
Total							

Grand Total 160 100

TOTAL UNDISTURBED TYPES (PERCENT) 15

Total Steps ea. CT. _____
----- = Composition
Grand Total Steps _____

✓ 15 Status (check)
 _____ 0 - 15 = very early seral
 _____ 16 - 40 = early seral
 _____ 41 - 60 = mid seral
 _____ 61 - 85 = late seral
 _____ 85 + = PNC

FOR MORE INFORMATION

You'll find detailed information on many aspects of rangeland and riparian management in these titles and in other publications, slide sets, CD-ROMs, and videos from UC ANR:

California Guidelines for Residual Dry Matter (RDM) Management on Coastal and Foothill Annual Rangelands, publication 8092

Sediment Delivery Inventory and Monitoring: A Method for Water Quality Management in Rangeland Watersheds, publication 8014

Visual Assessment of Riparian Health, publication 8089

To order these products, visit our online catalog at <http://anrcatalog.ucdavis.edu>. You can also place orders by mail, phone, or FAX, or request a printed catalog of publications, slide sets, CD-ROMs, and videos from

University of California
Agriculture and Natural Resources
Communication Services
6701 San Pablo Avenue, 2nd Floor
Oakland, California 94608-1239

Telephone: (800) 994-8849 or (510) 642-2431, FAX: (510) 643-5470

e-mail inquiries: danrcs@ucdavis.edu

An electronic version of this publication is available on the ANR Communication Services website at <http://anrcatalog.ucdavis.edu>

Publication 8094

© 2003 by the Regents of the University of California, Division of Agriculture and Natural Resources. All rights reserved.

The University of California prohibits discrimination against or harassment of any person employed by or seeking employment with the University on the basis of race, color, national origin, religion, sex, physical or mental disability, medical condition (cancer-related or genetic characteristics), ancestry, marital status, age, sexual orientation, citizenship, or status as a covered veteran (special disabled veteran, Vietnam-era veteran or any other veteran who served on active duty during a war or in a campaign or expedition for which a campaign badge has been authorized).

University Policy is intended to be consistent with the provisions of applicable State and Federal laws.

Inquiries regarding the University's nondiscrimination policies may be directed to the Affirmative Action/Staff Personnel Services Director, University of California, Agriculture and Natural Resources, 300 Lakeside Drive, 6th Floor, Oakland, CA 94612-3550 (510) 987-0096. **For information about obtaining this publication, call (800) 994-8849. For downloading information, call (530) 754-5112.**

pr-12/03-WJC/VJG



This publication has been anonymously peer reviewed for technical accuracy by University of California scientists and other qualified professionals. This review process was managed by the ANR Associate Editor for Natural Resources.