
# Optimizing water productivity in food production

THE ROSENBERG FORUM, ZARAGOZA, 2008

**ELIAS FERERES** 

INSTITUTE FOR SUSTAINABLE AGRICULTURE-CSIC & UNIVERSITY OF CORDOBA, SPAIN




#### PRODUCTION PER UNIT WATER USED


Infrastructures (from the source to the irrigation system), institutions, markets, ....; genotypes, environmental conditions, and....




EFFICIENCY OF WATER USE IS OFTEN LIMITED BY FACTORS OTHER THAN WATER,

IMPROVING
MANAGEMENT AT
ALL LEVELS IS
MOST CRITICAL







China Loess Plateau

#### Improving WP in rainfed systems:

WP= Y/ET Options: increase Y or decrease ET

Yield= Transpiration x Transp. Eff. (WPb) X Harvest Index Y (kg/ha)= T (mm) x TE (kg/ha/mm) x HI (kg/kg)

Options: Biological, Environmental and Management

What can we expect from GM crops?
Can we predict/change the environment?
How to best manage rainfed systems?
Objectives: maximize the fraction of rainfall into T and maintain HI

### New Indian Bed Planter Planting Cotton on Permanent Beds in Uzbekistan



# Improving WP in Irrigated Agriculture What has been happening?

1. Changes in irrigation systems.

2. Improved ET estimates and scheduling

3. Shift from low value to high value crops.


## (Water saved has been used to expand irrigated areas)

## The challenges to further increase Wp in irrigated agriculture

- •How much water would be available this season?
- •How should I distribute it among crops?
- Optimizing scheduling under Deficit Irrigation
- Applying the water as uniform as possible
- Monitoring, evaluation, and real-time feedbacks to improve performance



### EFFECTS OF IRRIGATION UNIFORMITY ON THE RELATION BETWEEN YIELD AND IRRIGATION WATER



# Under Water scarcity, Reduce Water Consumption (ET)

Therefore decrease

Eand/or T

### **DEFICIT IRRIGATION = APPLICATIONS BELOW CROP WATER REQUIREMENTS (ET)**

(Soil water deficits may or may not cause crop water deficits and reduction in ET)

#### **REGULATED DEFICIT IRRIGATION (RDI) =**


PLANNED CROP WATER
DEFICITS AT SPECIFIC DEVELOPMENTAL STAGES.

(Crop water stress occurs at certain stages; ET may or may not be reduced significantly; RISKS increased)

## Five-year peach experiment in Cordoba, where RDI and SDI were compared to full irrigation; in both DI treatments, 66% of the full requirements was applied

				<b>发</b> 为(
Treatment	Yield (t ha <sup>-1</sup> )	Fruit volume (FV cm³)	Relative FV	
Full	42.2 a	213 a	100	
SDI	38.6 b	198 b	92.9	×
RDI	41.2 a	213 a	100	
			Soriano & Fe	reres


## There is always an optimal Deficit Irrigation strategy that must be determined through long-term research!






#### COTTON YIELD RESPONSE TO WATER IN SOUTHERN SPAIN

The yield-response function was generated for wet (R), normal (N), and dry (D) years, using a computer simulation model.




## **AquaCrop**: a simulation model of water-limited crop production, currently being developed in FAO by P. Steduto, T. Hsiao, D. Raes, and E. Fereres.





#### FARM INCOME AS A FUNCTION OF APPLIED WATER AND WATER COSTS

Cotton gross margin in Southern Spain under current CAP of the EU




# IF THE SUPPLY IS RESTRICTED BELOW NORMAL, WHAT WOULD BE THE OPTIMAL COMBINATION OF CROPS IN MY FARM?






#### Spain, RDI strategy





Water Allocation: 3000m<sup>3</sup>/ha (1ac/ft)

Income (FI) (€/ha)
Maize 919.3
Cotton 813.0
Wheat 520.1


#### **ASSESSING IRRIGATION PERFORMANCE**







Large variations in Crop ET among farms



ET (mm) 0.0

100

400

700 1000 Water Limitations of satellite imagery: Time interval (every 14 days), problems with thermal sensors, pixel size, etc..









P. Zarco's team, IAS, Córdoba.









Flights over a field of 30 maize cultivars replicated three times, showing images of canopy temperature diferential where variations among cultivars are detected prior to irrigation

