## UCCE/DWR Weekly Crop Water Use Report





## WEEKLY SOIL MOISTURE LOSS IN INCHES

(Estimated Crop Evapotranspiration or  $ET_C$ ) 06/02/23 through 06/08/23

Crops (Leafout Date)	;	#148 Merced				#39 Parlier			#258 Lemon Cove			
	6/2 - 6/8	Accum'd	6/9 - 6/15		6/2 - 6/8	Accum'd	6/9 - 6/15		6/2 - 6/8	Accum'd	6/9 - 6/15	
	Water	Seasonal	Estimated		Water	Seasonal	Estimated		Water	Seasonal	Estimated	
	Use	Water Use	ETc		Use	Water Use	ETc		Use	Water Use	ЕТс	
Almonds (3/13) *	1.56	13.40	1.87		1.45	14.23	1.87		1.44	13.54	1.87	
Pistachio (4/20) * **	1.60	6.91	2.06		1.50	7.25	2.06		1.48	7.00	2.06	
Citrus (2/1)	1.05	13.10	1.22		0.99	14.03	1.22		0.98	13.32	1.22	
Raisin Grapes (3/22) (11 ft. row spacing)	1.05	4.58	1.28		0.98	4.82	1.28		0.97	4.68	1.28	
Winegrapes (3/22) (10 ft. spacing on California Sprawl Trellis)	1.02	4.79	1.28		0.95	5.07	1.28		0.94	4.89	1.28	
Walnuts (4/20)	1.27	6.61	1.57		1.18	7.02	1.57		1.17	6.82	1.57	
Stone Fruit (3/19)	1.36	9.34	1.57		1.25	9.84	1.57		1.26	9.48	1.57	
Past 7 days precipitation (inches)		0.00			<u>-</u>	0.00	_			0.15	_	
Accumulated precipitation (inches) (1/1/2023)		16.25				12.53				15.72		

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

<sup>\*\*</sup> Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 – resulting in about 8% greater water use than shown in these tables.

DAGE WEEKING A DRI IER WARER IN INCHES	A DILICEPO POD PERIORIA I
PAST WEEKLY APPLIED WATER IN INCHES.	

Crops								-	#258 Lemon Cove				
	#140 Merceu					#39 Parlier							
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%	
Almonds (3/13)	2.4	2.1	1.8	1.6	2.2	1.9	1.7	1.5	2.2	1.9	1.7	1.5	
Pistachio (4/20)	2.5	2.1	1.9	1.7	2.3	2.0	1.8	1.6	2.3	2.0	1.7	1.6	
Citrus (2/1)	1.6	1.4	1.2	1.1	1.5	1.3	1.2	1.0	1.5	1.3	1.2	1.0	
Raisin Grapes (3/22) (11 ft. row spacing)	As	sume all gr	ape	1.1	Assume all grape 1.0			1.0	Assume all grape			1.0	
Winegrapes (3/22) (10 ft. spacing on California Sprawl Trellis)	irriga	ation type is	s drip	1.1	irrigation type is drip		1.0	irrigation type is drip			1.0		
Walnuts (4/20)	2.0	1.7	1.5	1.3	1.8	1.6	1.4	1.2	1.8	1.6	1.4	1.2	
Stone Fruit (3/19)	2.1	1.8	1.6	1.4	1.9	1.7	1.5	1.3	1.9	1.7	1.5	1.3	

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

DAOT WEETST V	ADDITED WATED	IN CALLON DED	TREE OR VINE
PASI WEEKLY	APPLIED WATER	IN GALLON PER	TREE OR VINE

Crops		ed			#39 Parlier	r		#258 Lemon Cove				
Almonds 115 Trees/A	567	496	425	378	519	449	401	354	519	449	401	354
Pistachio 106 Trees/A	623	523	473	424	573	498	448	399	573	498	424	399
Citrus 110 Trees/A	395	346	296	272	370	321	296	247	370	321	296	247
Raisin Grapes 566 Vines/A	Assume all grape		53	Assume all grape 48			48	Assume all grape			48	
Winegrapes 622 Vines/A	irriga	ation type is	drip	48	irrigation type is drip			44	irrigation type is drip			44
Walnuts 76 Trees/A	715	607	536	464	643	572	500	429	643	572	500	429
Stonefruit 172 Trees/A	332	284	253	221	300	268	237	205	300	268	237	205
For further information concerning all counties receiving this report, contact	the Fresno (	Co. Farm Ad	visor's offic	e at (559) 24	41-7526.							

<sup>\*</sup> Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

HEALTHY COMMUNITIES • HEALTHY CAL

## UCCE/DWR Weekly Crop Water Use Report





## WEEKLY SOIL MOISTURE LOSS IN INCHES

(Estimated Crop Evapotranspiration or  $ET_C$ ) 06/02/23 through 06/08/23

Crops (Leafout Date)	-	124 Panoch	0		#2 Five Points							
Crops (Learout Date)									†	#15 Stratfor		
	6/2 - 6/8	Accum'd	6/9-6/15		6/2 - 6/8	Accum'd	6/9- 6/15		6/2 - 6/8	Accum'd	6/9-6/15	1
	Water	Seasonal	Estimated		Water	Seasonal	Estimated		Water	Seasonal	Estimated	
	Use	Water Use	ETc		Use	Water Use	ETc		Use	Water Use	ETc	
Almonds (3/13) *	1.61	13.94	2.11		1.57	13.99	2.11		1.67	15.47	2.13	
Pistachio (4/20) * **	1.67	7.18	2.29		1.62	7.17	2.29		1.72	7.91	2.31	
Citrus (2/1)	1.07	13.59	1.38		1.06	13.80	1.38		1.13	15.13	1.40	
Raisin Grapes (3/22) (11 ft. row spacing)	1.05	4.41	1.41		1.03	4.49	1.41		1.09	4.86	1.43	
Winegrapes (3/22) (10 ft. spacing on California Sprawl Trellis)	1.01	4.67	1.38		0.98	4.73	1.38		1.05	5.16	1.39	
Walnuts (4/20)	1.33	6.86	1.80		1.28	6.83	1.80		1.35	7.61	1.82	
Stone Fruit (3/19)	1.41	9.79	1.82		1.38	9.85	1.82		1.47	10.87	1.84	
Past 7 days precipitation (inches)		0.06				0.00				0.00		
Accumulated precipitation (inches) (1/1/2023)		6.18				7.65				7.90		

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

<sup>\*\*</sup> Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 – resulting in about 8% greater water use than shown in these tables.

PA	PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY 1												
Crops	#124 Panoche					#2 Five Po	ints						
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%	
Almonds (3/13)	2.5	2.1	1.9	1.7	2.4	2.1	1.8	1.7	2.6	2.2	2.0	1.8	
Pistachio (4/20)	2.6	2.2	2.0	1.8	2.5	2.2	1.9	1.7	2.6	2.3	2.0	1.8	
Citrus (2/1)	1.6	1.4	1.3	1.1	1.6	1.4	1.2	1.1	1.7	1.5	1.3	1.2	
Raisin Grapes (3/22) (11 ft. row spacing)	As	sume all gr	ape	1.1	Assume all grape 1.1			1.1	Assume all grape			1.1	
Winegrapes (3/22) (10 ft. spacing on California Sprawl Trellis)	irriga	ation type is	s drip	1.1	irriga	ation type is	s drip	1.1	irrigation type is drip		s drip	1.1	
Walnuts (4/20)	2.0	1.8	1.6	1.4	2.0	1.7	1.5	1.3	2.1	1.8	1.6	1.4	
Stone Fruit (3/19)	2.2	1.9	1.7	1.5	2.1	1.8	1.6	1.5	2.3	2.0	1.7	1.5	

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

PAST WEEKLY APPLIEI	WATER IN GALLO	ON PER TREE OR VINE

Crops		#124 Panoc	he			<b>#2 Five Poi</b>	ints		#15 Stratford			
Almonds 115 Trees/A	590	496	449	401	567	496	425	401	614	519	472	425
Pistachio 106 Trees/A	648	548	498	448	623	548	473	424	648	573	498	448
Citrus 110 Trees/A	395	346	321	272	395	346	296	272	420	370	321	296
Raisin Grapes 566 Vines/A	Assume all grape			53	Ass	ssume all grape 53			Assume all grape			53
Winegrapes 622 Vines/A	irriga	tion type is	drip	48	irrigation type is drip 4			48	irriga	ation type is	s drip	48
Walnuts 76 Trees/A	715	643	572	500	715	607	536	464	750	643	572	500
Stonefruit 172 Trees/A	347	300	268	237	332	284	253	237	363	316	268	237
For further information concerning all counties receiving this report, contact	the Fresno (	Co. Farm Ad	visor's office	e at (559) 24	41-7526.							

<sup>\*</sup> Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.