

**WEEKLY SOIL MOISTURE LOSS IN INCHES**

(Estimated Crop Evapotranspiration or ET<sub>c</sub>)

05/24/24 through 05/30/24

Crops (Leafout Date)	#148 Merced			#39 Parlier			#258 Lemon Cove		
	05/24 - 05/30 Water Use	Accum'd Seasonal Water Use	05/31 - 06/06 Estimated ET <sub>c</sub>	05/24 - 05/30 Water Use	Accum'd Seasonal Water Use	05/31 - 06/06 Estimated ET <sub>c</sub>	05/24 - 05/30 Water Use	Accum'd Seasonal Water Use	05/31 - 06/06 Estimated ET <sub>c</sub>
Almonds (3/1) *	1.63	12.49	1.76	1.74	13.05	1.76	1.70	12.70	1.72
Pistachio (4/20) * **	1.39	4.10	1.77	1.48	4.37	1.77	1.44	4.27	1.73
Citrus (2/1)	1.13	12.19	1.19	1.23	12.83	1.19	1.21	12.48	1.16
Raisin Grapes (3/11) (11 ft. row spacing)	1.09	4.57	1.21	1.18	4.90	1.21	1.16	4.81	1.18
Winegrapes (3/11) (10 ft. spacing on California Sprawl Trellis)	1.05	4.78	1.19	1.13	5.05	1.19	1.11	4.97	1.16
Walnuts (4/20)	1.15	4.18	1.41	1.26	4.47	1.41	1.24	4.46	1.37
Stone Fruit (3/11)	1.27	6.86	1.40	1.36	7.28	1.40	1.32	7.14	1.37
<b>Past 7 days precipitation (inches)</b>	0.00			0.00			0.00		
<b>Accumulated precipitation (inches) (1/1/2024)</b>	14.98			8.98			9.71		

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

\* Estimates are for orchard floor conditions where vegetation is managed by some combination of atrip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

\*\* Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 – resulting in about 8% greater water use than shown in these tables.

**PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY <sup>1</sup>**

Crops	#148 Merced				#39 Parlier				#258 Lemon Cove			
	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
Almonds (3/1)	2.5	2.2	1.9	1.7	2.7	2.3	2.0	1.8	2.6	2.3	2.0	1.8
Pistachio (4/20)	2.1	1.9	1.6	1.5	2.3	2.0	1.7	1.6	2.2	1.9	1.7	1.5
Citrus (2/1)	1.7	1.5	1.3	1.2	1.9	1.6	1.4	1.3	1.9	1.6	1.4	1.3
Raisin Grapes (3/11) (11 ft. row spacing)	Assume all grape			1.1	Assume all grape			1.2	Assume all grape			1.2
Winegrapes (3/11) (10 ft. spacing on California Sprawl Trellis)	irrigation type is drip			1.1	irrigation type is drip			1.2	irrigation type is drip			1.2
Walnuts (4/20)	1.8	1.5	1.4	1.2	1.9	1.7	1.5	1.3	1.9	1.7	1.5	1.3
Stone Fruit (3/11)	2.0	1.7	1.5	1.3	2.1	1.8	1.6	1.4	2.0	1.8	1.6	1.4

<sup>1</sup> The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

**PAST WEEKLY APPLIED WATER IN GALLON PER TREE OR VINE**

Crops	#148 Merced				#39 Parlier				#258 Lemon Cove			
	Almonds 115 Trees/A	590	519	449	401	638	543	472	425	614	543	472
Pistachio 106 Trees/A	523	473	399	374	573	498	424	399	548	473	424	374
Citrus 110 Trees/A	420	370	321	296	469	395	346	321	469	395	346	321
Raisin Grapes 566 Vines/A	Assume all grape			53	Assume all grape			58	Assume all grape			58
Winegrapes 622 Vines/A	irrigation type is drip			48	irrigation type is drip			52	irrigation type is drip			52
Walnuts 76 Trees/A	643	536	500	429	679	607	536	464	679	607	536	464
Stonefruit 172 Trees/A	316	268	237	205	332	284	253	221	316	284	253	221

For further information concerning all counties receiving this report, contact the Fresno Co. Farm Advisor's office at (559) 241-7526.

**WEEKLY SOIL MOISTURE LOSS IN INCHES**

(Estimated Crop Evapotranspiration or ET<sub>c</sub>)

05/24/24 through 05/30/24

Crops (Leafout Date)	#124 Panoche			#2 Five Points			#15 Stratford		
	05/24- 05/30 Water Use	Accum'd Seasonal Water Use	05/31- 06/06 Estimated ETc	5/26 - 6/1 Water Use	Accum'd Seasonal Water Use	6/2 - 6/8 Estimated ETc	5/26 - 6/1 Water Use	Accum'd Seasonal Water Use	6/2 - 6/8 Estimated ETc
Almonds (3/1) *	1.84	13.17	1.88	1.92	13.57	1.93	1.60	12.27	1.94
Pistachio (4/20) * **	1.57	4.67	1.95	1.64	4.86	2.00	1.37	4.13	2.01
Citrus (2/1)	1.28	13.25	1.27	1.34	13.77	1.30	1.13	12.67	1.31
Raisin Grapes (3/11) (11 ft. row spacing)	1.23	5.07	1.31	1.30	5.28	1.34	1.07	4.59	1.35
Winegrapes (3/11) (10 ft. spacing on California Sprawl Trellis)	1.16	5.20	1.28	1.22	5.37	1.30	1.01	4.74	1.31
Walnuts (4/20)	1.33	4.72	1.53	1.39	4.93	1.58	1.17	4.28	1.59
Stone Fruit (3/11)	1.44	7.69	1.53	1.49	7.84	1.58	1.26	6.97	1.59
<b>Past 7 days precipitation (inches)</b>	0.00			0.00			0.00		
<b>Accumulated precipitation (inches) (1/1/2024)</b>	6.60			6.86			5.43		

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

\* Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

\*\* Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 – resulting in about 8% greater water use than shown in these tables.

**PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY <sup>1</sup>**

Crops	#124 Panoche				#2 Five Points				#15 Stratford			
	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
System Efficiency >>												
Almonds (3/1)	2.8	2.5	2.2	1.9	3.0	2.6	2.3	2.0	2.5	2.1	1.9	1.7
Pistachio (4/20)	2.4	2.1	1.8	1.7	2.5	2.2	1.9	1.7	2.1	1.8	1.6	1.4
Citrus (2/1)	2.0	1.7	1.5	1.3	2.1	1.8	1.6	1.4	1.7	1.5	1.3	1.2
Raisin Grapes (3/11) (11 ft. row spacing)	Assume all grape			1.3	Assume all grape			1.4	Assume all grape			1.1
Winegrapes (3/11) (10 ft. spacing on California Sprawl Trellis)	irrigation type is drip			1.2	irrigation type is drip			1.2	irrigation type is drip			1.1
Walnuts (4/20)	2.0	1.8	1.6	1.4	2.1	1.9	1.6	1.5	1.8	1.6	1.4	1.2
Stone Fruit (3/11)	2.2	1.9	1.7	1.5	2.3	2.0	1.8	1.6	1.9	1.7	1.5	1.3

<sup>1</sup> The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

**PAST WEEKLY APPLIED WATER IN GALLON PER TREE OR VINE**

Crops	#124 Panoche				#2 Five Points				#15 Stratford			
Almonds 115 Trees/A	661	590	519	449	708	614	543	472	590	496	449	401
Pistachio 106 Trees/A	598	523	448	424	623	548	473	424	523	448	399	349
Citrus 110 Trees/A	494	420	370	321	518	444	395	346	420	370	321	296
Raisin Grapes 566 Vines/A	Assume all grape			62	Assume all grape			67	Assume all grape			53
Winegrapes 622 Vines/A	irrigation type is drip			52	irrigation type is drip			52	irrigation type is drip			48
Walnuts 76 Trees/A	715	643	572	500	750	679	572	536	643	572	500	429
Stonefruit 172 Trees/A	347	300	268	237	363	316	284	253	300	268	237	205

For further information concerning all counties receiving this report, contact the Fresno Co. Farm Advisor's office at (559) 241-7526.