### University of California Agriculture and Natural Resources



## UCCE/DWR Weekly Crop Water Use Report

Making a Difference for California

#### WEEKLY SOIL MOISTURE LOSS IN INCHES

(Estimated Crop Evapotranspiration or  $ET_C$ ) 07/12/24 through 07/18/24

Crops (Leafout Date)	#	148 Merce	d		#39 Parlier			#258 Lemon Cove		
	07/12 - 07/18	Accum'd	07/19 - 07/25		07/12 - 07/18	Accum'd	07/19 - 07/25	07/12 - 07/18	Accum'd	07/19 - 07/25
	Water	Seasonal	Estimated		Water	Seasonal	Estimated	Water	Seasonal	Estimated
	Use	Water Use	ETc		Use	Water Use	ETc	Use	Water Use	ETc
Almonds (3/1) *	2.06	26.31	2.03		2.02	27.63	1.96	1.97	26.89	1.96
Pistachio (4/20) * **	2.13	18.59	2.10		2.07	19.59	2.03	2.03	19.12	2.03
Citrus (2/1)	1.26	20.86	1.26		1.23	22.03	1.19	1.20	21.41	1.19
Raisin Grapes (3/11) (11 ft. row spacing)	1.39	14.06	1.37		1.34	14.79	1.31	1.32	14.44	1.31
Winegrapes (3/11) (10 ft. spacing on California Sprawl Trellis)	1.57	15.11	1.55		1.54	15.93	1.49	1.50	15.55	1.49
Walnuts (4/20)	1.96	16.12	2.17		1.92	17.11	2.10	1.87	16.71	2.10
Stone Fruit (3/11)	1.99	18.66	1.96		1.95	19.81	1.89	1.90	19.29	1.89
Past 7 days precipitation (inches)		0.00		_		0.00			0.00	_
Accumulated precipitation (inches) (1/1/2024)		14.98				8.98			9.71	

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

### PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY 1

Crops		#148 Merce	ed			#39 Parlier			#258 Lemon Cove			
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
Almonds (3/1)	3.2	2.7	2.4	2.2	3.1	2.7	2.4	2.1	3.0	2.6	2.3	2.1
Pistachio (4/20)	3.3	2.8	2.5	2.2	3.2	2.8	2.4	2.2	3.1	2.7	2.4	2.1
Citrus (2/1)	1.9	1.7	1.5	1.3	1.9	1.6	1.4	1.3	1.8	1.6	1.4	1.3
Raisin Grapes (3/11) (11 ft. row spacing)	As	sume all gra	ipe	1.5	Assume all grape 1.4			1.4	Assume all grape			1.4
Winegrapes (3/11) (10 ft. spacing on California Sprawl Trellis)	irrig	ation type is	drip	1.7	irrigation type is drip		1.6	irrigation type is drip		1.6		
Walnuts (4/20)	3.0	2.6	2.3	2.1	3.0	2.6	2.3	2.0	2.9	2.5	2.2	2.0
Stone Fruit (3/11)	3.1	2.7	2.3	2.1	3.0	2.6	2.3	2.1	2.9	2.5	2.2	2.0

The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

#### PAST WEEKLY APPLIED WATER IN GALLON PER TREE OR VINE

Crops		#148 Merce	ed			#39 Parlier			#258 Lemon Cove			
Almonds 115 Trees/A	756	638	567	519	732	638	567	496	708	614	543	496
Pistachio 106 Trees/A	822	698	623	548	797	698	598	548	772	673	598	523
Citrus 110 Trees/A	469	420	370	321	469	395	346	321	444	395	346	321
Raisin Grapes 566 Vines/A	Assume all grape			72	Assume all grape 67			67	Assume all grape			67
Winegrapes 622 Vines/A	irrig	ation type is	drip	74	irrigation type is drip 70			70	irrigation type is drip			70
Walnuts 76 Trees/A	1072	929	822	750	1072	929	822	715	1036	893	786	715
Stonefruit 172 Trees/A	489	426	363	332	474	410	363	332	458	395	347	316

For further information concerning all counties receiving this report, contact the Fresno Co. Farm Advisor's office at (559) 241-7526.

<sup>\*</sup> Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

<sup>\*\*</sup> Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 – resulting in about 8% greater water use than shown in these tables.

# University of California Agriculture and Natural Resources Making a Difference for California



# UCCE/DWR Weekly Crop Water Use Report

### WEEKLY SOIL MOISTURE LOSS IN INCHES

(Estimated Crop Evapotranspiration or  $ET_C$ ) 07/12/24 through 07/18/24

Crops (Leafout Date)	#124 Panoche				#2 Five Points				#	d		
	07/12- 07/18	Accum'd	07/19- 07/25		07/12- 07/18	Accum'd	07/19- 07/25		07/12- 07/18	Accum'd	07/19- 07/25	
	Water	Seasonal	Estimated		Water	Seasonal	Estimated		Water	Seasonal	Estimated	
	Use	Water Use	ETc		Use	Water Use	ETc		Use	Water Use	ETc	
Almonds (3/1) *	2.05	28.20	2.05		2.20	29.17	2.17		1.89	26.14	2.17	
Pistachio (4/20) * **	2.12	20.41	2.12		2.26	21.16	2.24		1.92	18.68	2.24	
Citrus (2/1)	1.26	22.72	1.26		1.35	23.60	1.33		1.14	21.38	1.33	
Raisin Grapes (3/11) (11 ft. row spacing)	1.37	15.30	1.39		1.46	15.90	1.43		1.24	14.05	1.43	
Winegrapes (3/11) (10 ft. spacing on California Sprawl Trellis)	1.57	16.41	1.56		1.68	16.98	1.67		1.43	15.03	1.67	
Walnuts (4/20)	1.96	17.79	2.19		2.09	18.49	2.31		1.79	16.35	2.31	
Stone Fruit (3/11)	2.01	20.67	1.98		2.17	21.34	2.10		1.82	18.93	2.10	
Past 7 days precipitation (inches)		0.00				0.00		-		0.00	<del>-</del>	
Accumulated precipitation (inches) (1/1/2024)		6.60				6.86				5.43		

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

<sup>\*\*</sup> Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 – resulting in about 8% greater water use than shown in these tables.

PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY 1												
Crops	#124 Panoche					#2 Five Poi	nts		#15 Stratford			
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
Almonds (3/1)	3.2	2.7	2.4	2.2	3.4	2.9	2.6	2.3	2.9	2.5	2.2	2.0
Pistachio (4/20)	3.3	2.8	2.5	2.2	3.5	3.0	2.7	2.4	3.0	2.6	2.3	2.0
Citrus (2/1)	1.9	1.7	1.5	1.3	2.1	1.8	1.6	1.4	1.8	1.5	1.3	1.2
Raisin Grapes (3/11) (11 ft. row spacing)	As	ssume all gra	ipe	1.4	Assume all grape 1.5			1.5	Assume all grape			1.3
Winegrapes (3/11) (10 ft. spacing on California Sprawl Trellis)	irrig	ation type is	drip	1.7	irrigation type is drip			1.8	irrigation type is drip			1.5
Walnuts (4/20)	3.0	2.6	2.3	2.1	3.2	2.8	2.5	2.2	2.8	2.4	2.1	1.9
Stone Fruit (3/11)	3.1	2.7	2.4	2.1	3.3	2.9	2.6	2.3	2.8	2.4	2.1	1.9

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

DACT WEELL V	ADDITED WATED	IN CALLON DED	TOPE OF UNIT
PASI WHEELY	APPLIED WATER	IN CALLON PER	TRABLUK VINH.

	17151	WEEKLI	MI LILD W	TILLITING	TILLON I EI	K TREE OR	VIIVE					
Crops	#124 Panoche				#2 Five Poi	nts						
Almonds 115 Trees/A	756	638	567	519	803	685	614	543	685	590	519	472
Pistachio 106 Trees/A	822	698	623	548	872	747	673	598	747	648	573	498
Citrus 110 Trees/A	469	420	370	321	518	444	395	346	444	370	321	296
Raisin Grapes 566 Vines/A	Assume all grape			67	Assume all grape 72			72	Assume all grape			62
Winegrapes 622 Vines/A	irrig	ation type is	drip	74	irrigation type is drip 79			79	irrigation type is drip			65
Walnuts 76 Trees/A	1072	929	822	750	1143	1000	893	786	1000	857	750	679
Stonefruit 172 Trees/A	489	426	379	332	521	458	410	363	442	379	332	300
For further information concerning all counties receiving this report, contact	the Fresno C	o. Farm Advi	isor's office a	t (559) 241-7	7526.							

<sup>\*</sup> Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.