University of California Agriculture and Natural Resources Making a Difference for California



UCCE/DWR Weekly Crop Water Use Report

WEEKLY SOIL MOISTURE LOSS IN INCHES

(Estimated Crop Evapotranspiration or ET_C) 06/13/25 through 06/19/25

Crops (Leafout Date)	#148 Merced			#39 Parlier				#258 Lemon Cove				
	06/13 - 06/19	Accum'd	06/20 - 06/26	06/13 - 06/19	Accum'd	06/20 - 06/26		06/13 - 06/19	Accum'd	06/20 - 06/26		
	Water	Seasonal	Estimated	Water	Seasonal	Estimated		Water	Seasonal	Estimated		
	Use	Water Use	ETc	Use	Water Use	ETc		Use	Water Use	ETc		
Almonds (3/1) *	1.93	17.74	1.98	2.11	18.88	1.96		2.02	17.54	1.96		
Pistachio (4/25) * **	2.08	9.81	2.12	2.26	10.79	2.10		2.17	10.18	2.10		
Citrus (2/1)	1.40	16.49	1.42	1.54	17.56	1.40		1.47	16.32	1.40		
Raisin Grapes (4/14) (11 ft. row spacing)	1.35	8.02	1.39	1.45	8.72	1.37		1.40	8.18	1.37		
Winegrapes (4/14) (10 ft. spacing on California Sprawl Trellis)	1.47	8.09	1.52	1.60	8.84	1.50		1.54	8.32	1.50		
Walnuts (4/14)	1.59	9.46	1.77	1.76	10.38	1.75		1.68	9.76	1.75		
Stone Fruit (3/8)	1.59	11.60	1.71	1.76	12.63	1.69		1.68	11.77	1.69		
Past 7 days precipitation (inches)		0.00			0.00		_		0.00			
Accumulated precipitation (inches) (1/1/2025)		0.00			5.35				4.44			

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

* Estimates are for orchard floor conditions where vegetation is managed by some combination of sarip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

^{**} Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 - resulting in about 8% greater water use than shown in these tables.

DACT WEELL VADDITED	WATED IN INCH	DO ADHIOTED DOI	D DEDICIENCY 1
PAST WEEKLY APPLIED) WAIEK IN INCH	ES. ADJUSTED FO	K EFFICIENCY :

Crops		#148 Merce	d			#39 Parlier			#258 Lemon Cove			
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
Almonds (3/1)	3.0	2.6	2.3	2.0	3.2	2.8	2.5	2.2	3.1	2.7	2.4	2.1
Pistachio (4/25)	3.2	2.8	2.4	2.2	3.5	3.0	2.7	2.4	3.3	2.9	2.6	2.3
Citrus (2/1)	2.2	1.9	1.6	1.5	2.4	2.1	1.8	1.6	2.3	2.0	1.7	1.5
Raisin Grapes (4/14) (11 ft. row spacing)	As	sume all gra	pe	1.4	Assume all grape 1.5			1.5	A	1.5		
Winegrapes (4/14) (10 ft. spacing on California Sprawl Trellis)	irrig	ation type is	drip	1.5	irrigation type is drip		1.7	irrigation type is drip		drip	1.6	
Walnuts (4/14)	2.4	2.1	1.9	1.7	2.7	2.3	2.1	1.9	2.6	2.2	2.0	1.8
Stone Fruit (3/8)	2.4	2.1	1.9	1.7	2.7	2.3	2.1	1.9	2.6	2.2	2.0	1.8

¹ The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

PAST WEEKLY APPLIED WATER IN GALLON PER TREE OR VINE

	1110	I III	THE PERSON	TITE IN C	TELECTION	· III	7 21 123					
Crops		#148 Merce	ed			#39 Parlier			#258 Lemon Cove			
Almonds 115 Trees/A	708	614	543	472	756	661	590	519	732	638	567	496
Pistachio 106 Trees/A	797	797 698 598		548	872	747	673	598	822	722	648	573
Citrus 110 Trees/A	543	469	395	370	592	518	444	395	568	494	420	370
Raisin Grapes 566 Vines/A	A	ssume all gra	pe	67	Assume all grape 72			72	Assume all grape 7.			72
Winegrapes 622 Vines/A	irrig	ation type is	drip	65	irrigation type is drip 74			irrigation type is drip			70	
Walnuts 76 Trees/A	857	750	679	607	965	822	750	679	929	786	715	643
Stonefruit 172 Trees/A	379	332	300	268	426	363	332	300	410	347	316	284
For further information concerning all counties receiving this report, contact	the Fresno Co	. Farm Advis	or's office at	(559) 241-75	26.							

University of California Agriculture and Natural Resources Making a Difference for California



UCCE/DWR Weekly Crop Water Use Report

WEEKLY SOIL MOISTURE LOSS IN INCHES

(Estimated Crop Evapotranspiration or ET_C) 06/13/25 through 06/19/25

Crops (Leafout Date)	#	124 Panoch	ie	#	2 Five Point	ts	#15 Stratford				
	06/13- 06/19	Accum'd	06/20- 06/26	06/13- 06/19	Accum'd	06/20- 06/26		06/13- 06/19	Accum'd	06/20- 06/26	
	Water	Seasonal	Estimated	Water	Seasonal	Estimated		Water	Seasonal	Estimated	
	Use	Water Use	ETc	Use	Water Use	ETc		Use	Water Use	ETc	
Almonds (3/1) *	2.18	18.70	2.19	2.33	19.43	2.23		2.36	20.60	2.23	
Pistachio (4/25) * **	2.34	10.88	2.29	2.49	11.39	2.31		2.53	11.94	2.31	
Citrus (2/1)	1.60	17.96	1.59	1.71	18.55	1.61		1.72	19.73	1.61	
Raisin Grapes (4/14) (11 ft. row spacing)	1.50	8.81	1.53	1.61	9.25	1.55		1.64	9.75	1.55	
Winegrapes (4/14) (10 ft. spacing on California Sprawl Trellis)	1.62	8.88	1.65	1.75	9.27	1.67		1.78	9.80	1.67	
Walnuts (4/14)	1.83	10.54	1.94	1.96	10.96	1.96		1.99	11.54	1.96	
Stone Fruit (3/8)	1.83	12.72	1.88	1.96	13.14	1.90		1.99	13.83	1.90	
Past 7 days precipitation (inches)		0.00			0.00		•		0.00		
Accumulated precipitation (inches) (1/1/2025)		2.31			3.13				2.72		

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

^{**} Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 - resulting in about 8% greater water use than shown in these tables.

	PAST WEI	EKLY APPI	JED WATE	R IN INCHE	ES, ADJUSTE	D FOR EFF	ICIENCY 1					
Crops		#124 Panocl	he			#2 Five Poi	nts					
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
Almonds (3/1)	3.4	2.9	2.6	2.3	3.6	3.1	2.7	2.5	3.6	3.1	2.8	2.5
Pistachio (4/25)	3.6	3.1	2.8	2.5	3.8	3.3	2.9	2.6	3.9	3.4	3.0	2.7
Citrus (2/1)	2.5	2.1	1.9	1.7	2.6	2.3	2.0	1.8	2.6	2.3	2.0	1.8
Raisin Grapes (4/14) (11 ft. row spacing)	As	ssume all gra	ipe	1.6	Assume all grape 1.7			1.7	As	1.7		
Winegrapes (4/14) (10 ft. spacing on California Sprawl Trellis)	irrig	gation type is	drip	1.7	irrig	irrigation type is drip			irrigation type is drip			1.9
Walnuts (4/14)	2.8	2.4	2.2	1.9	3.0	2.6	2.3	2.1	3.1	2.7	2.3	2.1
Stone Fruit (3/8)	2.8	2.4	2.2	1.9	3.0	2.6	2.3	2.1	3.1	2.7	2.3	2.1

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

	PAST	T WEEKLY	APPLIED W	VATER IN G	GALLON PER	TREE OR	VINE					
Crops		#124 Panoc	he			#2 Five Poi	nts					
Almonds 115 Trees/A	803	685	614	543	850	732	638	590	850	732	661	590
Pistachio 106 Trees/A	897	772	698	623	947	822	722	648	972	847	747	673
Citrus 110 Trees/A	617	518	469	420	642	568	494	444	642	568	494	444
Raisin Grapes 566 Vines/A	A	ssume all gra	ipe	77	Assume all grape 82			Assume all grape			82	
Winegrapes 622 Vines/A	irrig	gation type is	drip	74	irrig	irrigation type is drip			irrigation type is drip			83
Walnuts 76 Trees/A	1000	857	786	679	1072	929	822	750	1108	965	822	750
Stonefruit 172 Trees/A	442	379	347	300	474	410	363	332	489	426	363	332
For further information concerning all counties receiving this report, contact	t the Fresno C	o. Farm Advi	sor's office at	(559) 241-7	526.							

^{*} Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.